

Preoperative Evaluation of Lymph Node Metastasis in Esophageal Cancer

Yoko Murata, Masaho Ohta, Kazuhiko Hayashi, Hiroko Ide, and Ken Takasaki

Lymph node metastasis (LMN) in esophageal cancer occurs from the superficial cancer and spreads wildly from the neck to the abdomen. Hence precise determination of LMN is essential when adequate treatments are employed. There are several reports about ultrasonic features of malignant lymph nodes (LNs), and in summary, reports of endoscopic ultrasonography (EUS) findings of malignant LNs showed they were more than 5-10 mm in diameter, with a distinct border, hypoechoic internal echo and round shape. Sensitivity, specificity and accuracies for the diagnosis of malignant LNs by EUS were 49-99%, 33-99% and 71-96%. The rates widely varied, because the accuracy of EUS's ability to determine malignancy were based on the evaluation of various echo features of LNs, and were dependent on the judgement of subjective observers. Therefore histological analysis is necessary for adequate treatments. Endoscopic ultrasonography guided fine-needle aspiration cytology (EUS-FNA) has been performed for the diagnosis of malignant LNs since 10 years. Results of those reports were sensitivity 81-97%, specificity 83-100% and accuracy 83-97%. EUS-FNA staging was better than EUS staging. Also clinically obvious complications by EUS-FNA have not been reported. Therefore published evidence showed that EUS-FNA is safe and useful for confirmation of malignant LNs. (Ann Thorac Cardiovasc Surg 2003; 9: 88-92)

Key words: endoscopic ultrasonography (EUS), endoscopic ultrasonography guided fine-needle aspiration cytology (EUS-FNA), diagnosis of lymph node metastasis (LNM)

Introduction

Lymph node metastasis (LNM) in esophageal cancer occurs from the superficial cancer and spreads from the neck, the mediastinum and the abdomen. There are several papers that have described the high incidence of LNM in esophageal cancer. The Japanese Society for Esophageal Diseases reported that around 60% of cases have LNM. Kajiyama et al.¹⁾ reported that frequency of LNM in esophageal cancer was 22% in the neck, 34% in the upper mediastinum, 23% in midmediastinum, 20% in the lower mediastinum, and 38% in the upper area of the abdomen. Shiozaki et al.²⁾ reported that 46% of pT1 tumors have

mediastinal lymph node (LN) involvement, 42% for the recurrent nerve chain nodes, 21% for the cervical nodes, 26% for the abdominal nodes, pT2 tumors have 53%, 40%, 20%, 33%, pT3 tumors have 73%, 43%, 28%, 60%, pT4 tumors have 83%, 58%, 50%, 75%, respectively. As the depth of tumor invasion is deeper, the frequency of nodal involvement increases. As to the tumor location, the rate of LNM of the upper thoracic esophageal cancer was higher in cervical and thoracic LN, the rate of LNM of middle thoracic esophageal cancer was wide spread from the cervical LN to the abdominal LN, the rate of LNM of lower thoracic esophageal cancer was higher in the abdominal LN and the thoracic LN. Hence the rate of LNM is influenced by tumor location but there is an overlap. Therefore precise determination of LNM in esophageal cancer is important for selection of adequate treatments. Several modalities, such as computed tomography (CT), magnetic resonance imaging (MRI), ultrasonography, endoscopic ultrasonography (EUS), positron emission tomography (PET) have been employed for di-

From Department of Surgery I, Tokyo Women's Medical University, Tokyo, Japan

Received December 2, 2002; accepted for publication December 12, 2002.

Address reprint requests to Yoko Murata, Department of Surgery I, Tokyo Women's Medical University, 8-1 Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan.

Table 1. EUS findings of nonmetastatic and metastatic LN⁶⁾

	Nonmetastatic LN	Metastatic LN
Size (mm)		
Longest-D	8.5±5.4	10.8±6.5
Shortest-D	4.3±2.0	8.4±13.7
Morphology (%)		
Distinct border	24	76
Indistinct border	95	5
Heterogeneous internal echo	5	95
Homogeneous internal echo	90	10

EUS, endoscopic ultrasonography; LN, lymph node; D, diameter.

Table 2. Echo features of benign and malignant LN⁹⁾

Echo features	Benign (%)	Malignant (%)
Size (≥1 cm)	74	94
Distinct margins	26	50
Shape		
Round	16	31
Oval	53	44
Triangle	26	6
Central echogenic area	21	0
Hypoechoic nature	100	100
Mixed echogenicity	0	31

Table 3. Association between EUS features and LN histopathology¹⁰⁾

	Benign		Malignant	
Size of LN (mm)	1.4±0.1		1.7±0.1	
Morphology (score)	9.9±0.3		11.9±0.3	
Visual analog of LN				
Roundness	Irregular	Oval	Round	
Score	1	2	3	4
Echogenicity	Hyperechoic	Intermediate	Hypoechoic	
Score	1	2	3	4
Homogeneity	Heterogeneous	Intermediate	Homogeneous	
Score	1	2	3	4

agnosis of LNM, however, EUS provides higher accuracy among those modalities.³⁻⁵⁾ In this paper, the role of EUS for diagnosis of LNM has been reviewed.

EUS Feature of LNM

There are several papers describing malignant LNs. Murata et al.⁶⁾ reported the size and morphological EUS features of metastatic LNs. They were more than 5 mm in the shortest dimension, had a distinct border and heterogeneous hypoechoic internal echo (Table 1). Tio et al.⁷⁾ reported the criteria of LNM as hypoechoic clearly delineated nodes. Heintz et al.⁸⁾ described that the size of malignant LNs (10.1±5.3 mm) were significantly larger than benign nodes (7.4±2.9 mm). Comparing malignant and benign LNs,⁹⁾ in malignant nodes, there was a higher prevalence of nodes being larger than 1 cm, and having hypoechoic distinct margins and a round shape (Table 2). All visualized LNs were characterized according to standard EUS figures and scales were settled. Malignant LNs have higher scores than benign nodes (Table 3).¹⁰⁾ But the size of LNs between them was not much different. Summarizing those reports, malignant LNs should be

more than 5-10 mm, hypoechoic, have a distinct border and a round shape (Fig. 1). However, some EUS features overlapped between benign and malignant LNs so that there are no firmly established criteria.

Accuracy of Determination of LNM

The overall results of EUS of LNM are shown in Table 4. Sensitivities are from 49 to 99%, specificities are from 33 to 99% and accuracies are from 71 to 96%.^{6,7,11-20)} Misjudgements on EUS can be due to inadequate scanning, such as nontraversable tumors and insufficient penetration. Therefore facilities which report a high rate of advanced tumors may be carrying out less accurate EUS in which case CT is superior to EUS.³⁾ When the specificity increases, the sensitivity lessens, it might be influenced by established EUS features of malignant LN. On the other hand, microscopic infiltration can cause false negative as inflammatory swelling LNs cause false negative results. Accordingly the accuracy of EUS was different depending on the facilities condition, the judgment of various observers and overlap between features of benign and malignant LNs.




Fig. 1. EUS figure of malignant LN.
Right: the LN of resected specimen.

Table 4. The results of published data for accuracy of EUS in LNM

Authors	No. of patient	Sensitivity (%)	Specificity (%)	Accuracy (%)		
Murata et al. 1987 ⁶⁾	222	87	90	89		
Tio et al. 1990 ⁷⁾	113	99	50	84		
Fok et al. 1992 ¹¹⁾						
	CAN	98	58	76	71	
	Med		85	86		
	Abd	45	62	86		
Dittler et al. 1993 ¹²⁾	167	75	70	73		
Murata et al. 1994 ¹³⁾	328	85	91	90		
Binmoeller et al. 1995 ¹⁴⁾	38	90	44	79		
Natsugoe et al. 1996 ¹⁵⁾	37	80	88	T1	87	
Shimizu et al. 1997 ¹⁶⁾	82	49	99	95		
Vickers et al. 1998 ¹⁷⁾	49	97	54	86		
Nishimaki et al. 1999 ¹⁸⁾	224	78	60	72		
Salminen et al. 1999 ¹⁹⁾	32	95	33	72		
Catalano et al. 1999 ²⁰⁾				73		
	Med	149	79	63		
	CAN		83	98	FNA	96

CAN, celiac axis node; Med, mediastinum; Abd, abdomen; FNA, fine-needle aspiration cytology.

Endoscopic Ultrasonography Guided Fine-needle Aspiration Cytology (EUS-FNA)

Histological analysis of LNM is essential when indication for operation and LN dissection were determined or endoscopic treatments and less invasive treatments were introduced. For instance, when histological analysis was positive in the celiac node, operation could be avoided. Therefore video-assisted thoracoscopy and laparoscopy (VATS/LAP) were performed, and positive EUS results were 62% but VATS/LAP was 86%, 8 negative EUS re-

sults were confirmed and 6 positive by VATS/LAP.²¹⁾ The introduction of the curved linear array echoendoscope led to development of endoscopic ultrasonography guided fine-needle aspiration cytology (EUS-FNA) (Fig. 2). In 10 years, EUS-FNA has been introduced in the diagnosis of LNM. The decision for EUS-FNA was performed based on EUS appearance of the LN, such as suspected malignant or swelling. Adequate specimens for cytologic study were obtained in 90% of the procedures.²³⁾ EUS-FNA produced sensitivity 81-97%, specificity 83-100% and accuracy 83-97% (Table5).²²⁻²⁹⁾ And the accuracy of EUS-




Fig. 2. The curved linear array echoendoscope led to EUS-FNA.

FNA is higher than that of EUS.³⁰⁾ However, in patients whom have severe stenosis, the probe can not be passed and EUS-FNA is not possible, but dilation of the esophageal stenosis allows EUS-FNA to be performed, and accurate staging is possible.³¹⁾ There are no reports which describe severe complications caused by EUS-FNA. Furthermore, EUS-FNA may be a more cost-effective model than mediastinoscopy.³²⁾

In conclusion, EUS and EUS-FNA are safe and effective modalities for diagnosis of malignant LNs.

References

1. Kajiyama Y, Tsurumaru M. Esophagectomy with lymph node dissection through right thoracotomy. *Nippon Geka Gakkai Zasshi* 2002; **103**: 343–7.
2. Shiozaki H, Yano M, Tsujinaka T, et al. Lymph node metastasis along the recurrent nerve chain is an indication for cervical lymph node dissection in thoracic esophageal cancer. *Dis Esophagus*, 2001; **14**: 191–6.
3. Rösch T. Endosonographic staging of esophageal cancer: a review of literature results. *Gastrointest Endosc Clin N Am* 1995; **5**: 537–47.
4. Van Dam J. Endosonographic evaluation of patient with esophageal cancer. *Chest* 1997; **112**: 184S–190S.
5. Kulling D, Feldman DR, Kay CL, et al. Local staging of esophageal cancer using endoscopic magnetic resonance imaging: prospective comparison with endoscopic ultrasound. *Endoscopy* 1998; **30**: 745–9.
6. Murata Y, Muroi M, Yoshida M, Ide H, Hanyu F. Endoscopic ultrasonography in the diagnosis of esophageal carcinoma. *Surg Endosc* 1987; **1**: 11–6.
7. Tio TL, Coene PPLO, Luiken GJHM, Tytgat GJ. Endosonography in the clinical staging of esophagogastric carcinoma. *Gastrointest Endosc* 1990; **36**: S2–10.
8. Heintz A, Mildenerger P, Georg M, Garcia A, Junginger T. In vitro studies of lymph node analysis. *Gastrointest Endosc Clin N Am* 1995; **5**: 577–86.
9. Bhutani MS, Hawes RH, Hoffman BJ. A comparison of the accuracy of echo features during endoscopic ultrasound (EUS) and EUS-guided fine-needle aspiration for diagnosis of malignant lymph node invasion. *Gastrointest Endosc* 1997; **45**: 474–9.
10. Faige DO. EUS in patients with benign and malignant lymphadenopathy. *Gastrointest Endosc* 2001; **53**: 593–8.
11. Fok M, Cheng SWK, Wong J. Endosonography in patient selection for surgical treatment of esophageal carcinoma. *World J Surg* 1992; **16**: 1098–103.
12. Dittler HJ, Siewert JR. Role of endoscopic ultrasonography in esophageal carcinoma. *Endoscopy* 1993; **25**: 156–61.
13. Murata Y, Hayashi K, Kobayashi A, et al. Pre-operative staging of oesophageal carcinoma by ultrasound. *Asian J Surg* 1994; **17**: 200–7.
14. Binmoeller KF, Seifert H, Seitz U, Izbicki JR, Kida M, Soehendra N. Ultrasonic esophagoprobe for TNM staging of highly stenosing esophageal carcinoma. *Gastrointest Endosc* 1995; **41**: 547–52.
15. Natsugoe S, Yoshinaka H, Morinaga T, et al. Ultra-

Table 5. The results of published data for accuracy of EUS-FNA in LNM

Authors	No. of patient	Sensitivity (%)	Specificity (%)	Accuracy (%)
Giovannini et al. 1995 ²³⁾	24	81	100	83
Gress et al. 1997 ²⁴⁾	24	93	100	96
Hunerbein et al. 1998 ²⁵⁾	25	89	83	–
Silvestri et al. 1996 ²⁶⁾	26	89	100	92
Wiersema et al. 1998 ²⁷⁾	48	88	100	90
Williams et al. 1999 ²⁸⁾	60	87	100	90
Giovannini et al. 1999 ²⁹⁾	31	97	100	97

- sonographic detection of lymph-node metastases in superficial carcinoma of the esophagus. *Endoscopy* 1996; **28**: 674–9.
16. Shimizu Y, Mera K, Tsukagoshi H, et al. Endoscopic ultrasonography for the detection of lymph node metastasis in superficial esophageal carcinoma. *Dig Endosc* 1997; **9**: 178–82.
 17. Vickers J, Alderson D. Oesophageal cancer staging using endoscopic ultrasonography. *Br J Surg* 1998; **85**: 994–8.
 18. Nishimaki T, Tanaka O, Ando N, et al. Evaluation of accuracy of preoperative staging in thoracic esophageal cancer. *Ann Thorac Surg* 1999; **68**: 2059–64.
 19. Salminen JT, Farkkila MA, Ramo OJ, et al. Endoscopic ultrasonography in the preoperative staging of adenocarcinoma of the distal oesophagus and oesophagogastric junction. *Scand J Gastroenterol* 1999; **34**: 1178–82.
 20. Catalano MF, Alcocer E, Chak A, et al. Evaluation of metastatic celiac axis lymph nodes in patients with esophageal carcinoma: accuracy of EUS. *Gastrointest Endosc* 1999; **50**: 352–6.
 21. Luketich JD, Schauer P, Landreneau R, et al. Minimally invasive surgical staging is superior to endoscopic ultrasound in detecting lymph node metastases in esophageal cancer. *J Thorac Cardiovasc Surg* 1997; **114**: 817–23.
 22. Barawi M, Gress F. EUS-guided fine-needle aspiration in the mediastinum. *Gastrointest Endosc* 2000; **52** (Suppl): S12–7.
 23. Giovannini M, Seitz JF, Monges G, Perrier H, Rabbia I. Fine-needle aspiration cytology guided by endoscopic ultrasonography: results in 141 patients. *Endoscopy* 1995; **27**: 171–7.
 24. Gress F, Savides TJ, Sandler A, et al. Endoscopic ultrasonography, fine-needle aspiration biopsy guided by endoscopic ultrasonography, and computed tomography in the preoperative staging of non-small-cell lung cancer: a comparison study. *Ann Intern Med* 1997; **127**: 604–12.
 25. Hunerbein M, Ghadimi BM, Haensch W, Schlag PM. Transesophageal biopsy of mediastinal and pulmonary tumors by means of endoscopic ultrasound guidance. *J Thorac Cardiovasc Surg* 1998; **116**: 554–9.
 26. Silvestri GA, Hoffman BJ, Bhutani MS, et al. Endoscopic ultrasound with fine-needle aspiration in the diagnosis and staging of lung cancer. *Ann Thorac Surg* 1996; **61**: 1441–6.
 27. Wiersema MJ, Harada N, Daiehigh P, et al. Evaluation of mediastinal lymphadenopathy with transesophageal endosonography guided fine needle aspiration biopsy. *Acta Endoscopica* 1998; **28**: 7–19.
 28. Williams DB, Sahai AV, Aabakken L, et al. Endoscopic ultrasound guided fine needle aspiration biopsy: a large single centre experience. *Gut* 1999; **44**: 720–6.
 29. Giovannini M, Monges G, Seitz JF, et al. Distant lymph node metastasis in esophageal cancer: impact of endoscopic ultrasound-guided biopsy. *Endoscopy* 1999; **31**: 536–40.
 30. Vazquez-Sequeiros E, Norton ID, Clain JE, et al. Impact of EUS-guided fine-needle aspiration on lymph node staging in patients with esophageal carcinoma. *Gastrointest Endosc* 2001; **53**: 751–7.
 31. Wallace MB, Hawes RH, Sahai AV, Van Velse A, Hoffman BJ. Dilation of malignant esophageal stenosis to allow EUS guided fine-needle aspiration: safety on patient management. *Gastrointest Endosc* 2000; **51**: 309–13.
 32. Aabakken L, Silvestri GA, Hawes R, Reed CE, Marsi V, Hoffman B. Cost-efficacy of endoscopic ultrasonography with fine-needle aspiration vs. mediastinotomy in patients with lung cancer and suspected mediastinal adenopathy. *Endoscopy* 1999; **31**: 707–11.