

Surgical Treatment for Ischemic Mitral Regurgitation: Strategy for a Tethered Valve

Hiromu Terai, MD, Kouji Tao, MD, and Ryuzo Sakata, MD

Recently, ischemic mitral regurgitation (IMR) has been shown to be an individual risk factor for ischemic heart disease. The main mechanism of IMR is tethering of the leaflet secondary to left ventricular (LV) dilatation. In this situation, surgical treatment for IMR has been limited to ring annuloplasty with varying degrees of effectiveness. However, mid-term follow-up studies have shown that the results obtained with this approach are not satisfactory. Therefore, there has been a need to develop additional techniques to achieve more secure repair of IMR. The characteristics of the mitral leaflet configuration in IMR are apical displacement of the leaflets relative to the annulus, concavity of the leaflets, and a dilated annulus. Our basic strategy for a tethered mitral valve is rigid ring annuloplasty and inward correction of the outwardly displaced papillary muscle. For the latter correction, we employ the overlapping method or septal anterior ventricular exclusion (SAVE) procedure for LV volume reduction in cases of broad antero-septal infarction, or elevate the posterior papillary muscle by folding the LV wall at the root of the posterior papillary muscle via a small incision in the inferior wall in cases of infero-posterior infarction. An additional procedure is chordal cutting in combination with rigid ring annuloplasty and papillary muscle imbrication in combination with LV volume reduction. We have successfully combined these methods with the aid of detailed echocardiographic studies in individual patients. However, long-term follow-up will be necessary before this approach can be routinely adopted. (*Ann Thorac Cardiovasc Surg* 2005; 11: 288–92)

Key words: ischemic mitral regurgitation, tethering, surgical treatment

Introduction

Recently, attention is paid to ischemic mitral regurgitation (IMR) because it has become clear that IMR is an individual risk factor for ischemic heart disease.¹⁻⁵⁾ Previously, the main cause of IMR was thought to be mitral valve prolapse due to dysfunction of the papillary muscle.^{6,7)} However, the current opinion is that, in most cases, the main cause of IMR is tethering of the leaflet secondary to left

ventricular (LV) dilatation⁸⁻¹⁴⁾ and papillary muscle dysfunction attenuates IMR, as revealed by detailed echocardiographic studies.¹⁵⁾ In this situation, surgical treatment for IMR has been limited to annuloplasty using a rigid or flexible ring, with varying degrees of effectiveness.¹⁶⁻¹⁹⁾ However, Calafiore et al. have reported that ring annuloplasty is not effective in cases where tethering is strong,²⁰⁾ and recently McGee et al. reported that the short- and mid-term outcomes of ring annuloplasty are not satisfactory.²¹⁾ This may be because this procedure cannot address the tethering itself. Therefore, there has been a need to develop additional techniques to achieve more secure repair of IMR. Here we describe our surgical strategy for treatment of mitral valve tethering.

Characteristics of the mitral leaflet configuration in IMR
1: Apical displacement of the mitral leaflets relative to

From Department of Thoracic and Cardiovascular Surgery, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, Japan

Received June 22, 2005; accepted for publication July 6, 2005.
Address reprint requests to Hiromu Terai, MD: Department of Thoracic and Cardiovascular Surgery, Kagoshima University Graduate School of Medical and Dental Sciences, 8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan.

the annulus¹⁰⁻¹⁴⁾

This is caused by outward dislocation of the papillary muscle secondary to LV dilatation. It must be noted that in the case of infero-posterior infarction, this phenomenon may occur without such severe LV dilatation.²²⁻²⁵⁾

2: Concavity of the leaflet²⁶⁾

When there is heterogeneity of leaflet tethering, the tethering in the middle portion of the leaflet is stronger, resulting in concave deformation of the leaflet and narrowing of the coaptation zone.

3: Dilated annulus^{25,27-31)}

Dilatation of the annulus secondary to LV dilatation also plays a role in IMR. In this setting, the annulus is deformed, making it rounder and losing its physiological saddle shape.³²⁻³⁴⁾ The dilated distance of anterior and posterior annuli exacerbates the narrowed coaptation of the tethered valve.

Our strategy for a tethered mitral valve

Currently, no established method for IMR has been reported. In our institution, we combine the methods described below with the aid of detailed echocardiographic studies in individual patients.

Basic strategy

1: Ring annuloplasty

To increase the narrowed coaptation zone, this procedure is indispensable. If tethering is not so strong, this procedure alone achieves sufficient repair of IMR. To control the distance of the anterior and posterior annuli, we select a rigid ring.

2: Inward correction of the outwardly displaced papillary muscle

Considering the mechanism of IMR, the essential surgical aim must be inward correction of the displaced papillary muscle. For this purpose, we select the overlapping method^{35,36)} or septal anterior ventricular exclusion (SAVE) procedure^{37,38)} as the method of LV volume reduction in cases of broad antero-septal infarction accompanied by severe LV dilatation, or we elevate the posterior papillary muscle by folding the LV wall at the root of the posterior papillary muscle via a small incision in the inferior wall in cases of infero-posterior infarction (where the LV volume is often not increased to such an extent as to require reduction).³⁹⁾

Additional procedures

1: Chordal cutting^{40,41)}

When the shape of the leaflet is concave, this technique

is effective in gaining an adequate coaptation zone, and therefore we use it in combination with rigid ring annuloplasty.

2: Papillary muscle imbrication in combination with LV volume reduction⁴²⁾

When the distance of the anterior and posterior papillary muscles is too great for correction by LV volume reduction alone, we add this technique. It must be noted that this technique alone reduces LV volume.

Case presentations

*Case 1 (broad anterior infarction)*⁴³⁾

The concomitant operation is the left atrial Maze operation.

LV volume reduction was performed by the overlapping method.

In this case, although the degree of tethering was relatively small, the coaptation zone was markedly narrowed because of the concave shape. Four strut chordae of the anterior leaflet were cut, and then annuloplasty using a 30-mm Carpentier-Edwards rigid ring (Edwards Lifesciences, USA) was performed. After the operation, echocardiography showed a convex shape, an adequate coaptation zone, and absence of IMR (Fig. 1).

*Case 2 (broad anterior and inferior infarction)*³⁹⁾

The concomitant operation is CABG (lt.ITA-LAD · Ao-RCA).

LV volume reduction was also performed by the overlapping method.

In this case, two large posterior papillary muscles were present and both were strongly displaced by the presence of inferior infarction. Therefore, we imbricated the anterior one anteriorly and elevated the posterior one by folding the inferior wall. This eliminated most of the valve tethering when viewed through the left atriotomy before performing annuloplasty using a 28-mm Carpentier-Edwards rigid ring. After the operation, echocardiography showed an adequate coaptation zone and absence of tethering and IMR (Fig. 2).

Discussion

Detailed clinical studies using echocardiography, and also several experimental studies, conducted over the last few years have proved that the main mechanism responsible for IMR is outward displacement of the papillary muscle and consequent tethering of the mitral valve leaflets.^{1-5,8-15)} This means that IMR is not a disease of the valve, but of the LV myocardium, therefore, the surgical methods used




Fig. 1.

a. Case 1 before surgery.

The leaflets are apparently tethered towards the apex. The anterior leaflet shows concave deformity. Doppler mode demonstrates severe MR.

b. Case 1 after surgery.

MR and tethering of the leaflets have disappeared, and the leaflet is now the correct convex shape.




Fig. 2.

a. Case 2 before surgery.

The leaflets are apparently tethered towards the apex. The anterior leaflet shows slight concave deformity. Doppler mode demonstrates severe MR.

b. Case 2 after surgery.

MR and tethering of the leaflets have disappeared. However, concave deformity of the leaflet still remains.

previously for valvular MR are not effective except for ring annuloplasty.¹⁶⁻¹⁹⁾ However, recently a mid-term follow-up study showed that the outcome of ring annuloplasty

was not satisfactory.²¹⁾ Recently, several promising experimental trials have been reported. Tibayan et al. have reported that septal-lateral annular cinching i.e. reduc-

tion of the distance of the anterior and posterior annuli diminishes the severity of IMR,⁴⁴⁾ thus supporting our policy of selecting a rigid ring for annuloplasty. Messas et al. have reported a chordal cutting procedure,^{40,41)} and we have successfully adopted this method combined with rigid ring annuloplasty (case 1). Clinically, Hvass et al. have reported the use of a papillary muscle sling⁴⁵⁾ and Menicanti et al. have reported papillary muscle imbrication.⁴²⁾ These two methods share the same concept that reduction of the increased distance of the anterior and posterior papillary muscles diminishes the degree of tethering, with different approaches from the left atrium (LA) or the LV. We have also successfully adopted papillary muscle imbrication combined with LV volume reduction (case 2), and we consider that this method is indicated for cases of severe LV dilatation after broad anterior infarction.

Finally, a description of our original method is given below.³⁹⁾ To relocate the posterior papillary muscle inwardly, we fold the inferior LV wall at the root of the dislocated posterior papillary muscle. In combination with rigid ring annuloplasty, this method effectively diminishes IMR (case 2). We think that this approach is an essential one for reduction of tethering. However, its limitation is that the free wall of the LV at the root of the dislocated papillary muscle must be infarcted transmurally. Therefore, this method is good for cases of infero-posterior infarction, but not for severe LV dilatation after broad anterior infarction.

At present, irrespective of the method selected for treatment of IMR, the long-term results are unpredictable. Many factors can affect the outcome, including whether or not LV remodeling has been completed at the time of surgery, or whether it will continue after surgery. In any event, surgical therapy for IMR is still at the starting line, and long-term follow-up of many cases will be mandatory in order to establish the most suitable method.

References

1. Tchong JE, Jackman JD Jr, Nelson CL, et al. Outcome of patients sustaining acute ischemic mitral regurgitation during myocardial infarction. *Ann Intern Med* 1992; **117**: 18–24.
2. Lehmann KG, Francis CK, Dodge HT. Mitral regurgitation in early myocardial infarction. Incidence, clinical detection, and prognostic implications. TIMI Study Group. *Ann Intern Med* 1992; **117**: 10–7.
3. Lamas GA, Mitchell GF, Flaker GC, et al. Clinical significance of mitral regurgitation after acute myocardial infarction. Survival and Ventricular Enlargement Investigators. *Circulation* 1997; **96**: 827–33.
4. Grigioni F, Enriquez-Sarano M, Zehr KJ, Bailey KR, Tajik AJ. Ischemic mitral regurgitation: long-term outcome and prognostic implications with quantitative Doppler assessment. *Circulation* 2001; **103**: 1759–64.
5. Barzali B, Davis VG, Stone PH, Jaffe AS. Prognostic significance of mitral regurgitation in acute myocardial infarction. The MILIS Study Group. *Am J Cardiol* 1990; **65**: 1169–75.
6. Burch GE, DePasquale NP, Phillips JH. The syndrome of papillary muscle dysfunction. *Am Heart J* 1968; **75**: 399–415.
7. Hider CF, Taylor DE, Wade JD. The effect of papillary muscle damage on atrioventricular valve function in the left heart. *Q J Exp Physiol Cogn Med Sci* 1965; **50**: 15–22.
8. Ogawa S, Hubbard FE, Mardelli TJ, Dreifus LS. Cross-sectional echocardiographic spectrum of papillary muscle dysfunction. *Am Heart J* 1979; **97**: 312–21.
9. Godley RW, Wann LS, Rogers EW, Feigenbaum H, Weyman AE. Incomplete mitral leaflet closure in patients with papillary muscle dysfunction. *Circulation* 1981; **63**: 565–71.
10. Otsuji Y, Handschumacher MD, Schwammenthal E, et al. Insights from three-dimensional echocardiography into the mechanism of functional mitral regurgitation: direct vivo demonstration of altered leaflet tethering geometry. *Circulation* 1997; **96**: 1999–2008.
11. Yiu SF, Enriquez-Sarano M, Tribouilloy C, Seward JB, Tajik AJ. Determinants of the degree of functional mitral regurgitation in patients with systolic left ventricular dysfunction: A quantitative clinical study. *Circulation* 2000; **102**: 1400–6.
12. Kaul S, Spotnitz WD, Glasheen WP, Touchstone DA. Mechanism of ischemic mitral regurgitation. An experimental evaluation. *Circulation* 1991; **84**: 2167–80.
13. Dent JM, Spotnitz WD, Nolan SP, Jayaweera AR, Glasheen WP, Kaul S. Mechanism of mitral leaflet excursion. *Am J Physiol* 1995; **269** (6 pt 2): H2100–8.
14. He S, Fontaine AA, Schwammenthal E, Yoganathan AP, Levine RA. Integrated mechanism for functional mitral regurgitation: leaflet restriction versus coapting force: in vitro studies. *Circulation* 1997; **96**: 1826–34.
15. Messas E, Guerrero JL, Handschumacher MD, et al. Paradoxical decrease in ischemic mitral regurgitation with papillary muscle dysfunction: insights from three-dimensional and contrast echocardiography with strain rate measurement. *Circulation* 2001; **104**: 1952–7.
16. Bolling SF, Pagani FD, Deeb GM, Bach DS. Intermediate-term outcome of mitral reconstruction in cardiomyopathy. *J Thorac Cardiovasc Surg* 1998; **115**: 381–8.
17. Bach DS, Bolling SF. Improvement following correction of secondary mitral regurgitation in end-stage cardiomyopathy with mitral annuloplasty. *Am J Cardiol* 1996; **78**: 966–9.
18. Bach DS, Bolling SF. Early improvement in congestive heart failure after correction of secondary mitral

- regurgitation in end-stage cardiomyopathy. *Am Heart J* 1995; **129**: 1165–70.
19. Bax JJ, Braun J, Somer ST, et al. Restrictive annuloplasty and coronary revascularization in ischemic mitral regurgitation results in reverse left ventricular remodeling. *Circulation* 2004; **110** (11 Suppl 1): II103–8.
 20. Calafiore AM, Gallina S, Di Mauro M, et al. Mitral valve procedure in dilated cardiomyopathy: repair or replacement? *Ann Thorac Surg* 2001; **71**: 1146–52.
 21. McGee EC, Gillinov AM, Blackstone EH, et al. Recurrent mitral regurgitation after annuloplasty for functional ischemic mitral regurgitation. *J Thorac Cardiovasc Surg* 2004; **128**: 916–24.
 22. Izumi S, Miyatake K, Beppu S, et al. Mechanism of mitral regurgitation in patients with myocardial infarction: a study using real-time two-dimensional Doppler flow imaging and echocardiography. *Circulation* 1987; **76**: 777–85.
 23. Leor J, Feinberg MS, Vered Z, et al. Effect of thrombolytic therapy on the evolution of significant mitral regurgitation in patients with a first inferior myocardial infarction. *J Am Coll Cardiol* 1993; **21**: 1661–6.
 24. Gorman JH 3rd, Gorman RC, Plappert T, et al. Infarct size and location determine development of mitral regurgitation in the sheep model. *J Thorac Cardiovasc Surg* 1998; **115**: 615–22.
 25. Kumanohoso T, Otsuji Y, Yoshifuku S, et al. Mechanism of higher incidence of ischemic mitral regurgitation in patients with inferior myocardial infarction: quantitative analysis of left ventricular and mitral valve geometry in 103 patients with prior myocardial infarction. *J Thorac Cardiovasc Surg* 2003; **125**: 135–43.
 26. Nesta F, Otsuji Y, Handschumacher MD, et al. Leaflet concavity: a rapid visual clue to the presence and mechanism of functional mitral regurgitation. *J Am Soc Echocardiogr* 2003; **16**: 1301–8.
 27. Sabbah HN, Kono T, Rosman H, Jafri S, Stein PD, Goldstein S. Left ventricular shape: a factor in the etiology of functional mitral regurgitation in heart failure. *Am Heart J* 1992; **123**: 961–6.
 28. Ahmad RM, Gillinov AM, McCarthy PM, et al. Annular geometry and motion in human ischemic mitral regurgitation: novel assessment with three-dimensional echocardiography and computer reconstruction. *Ann Thorac Surg* 2004; **78**: 2063–8.
 29. Tibayan FA, Rodriguez F, Zasio MK, et al. Geometric distortions of the mitral valvular-ventricular complex in chronic ischemic mitral regurgitation. *Circulation* 2003; **108** (Suppl 1): II116–21.
 30. Green GR, Dagum P, Glasson JR, et al. Mitral annular dilatation and papillary muscle dislocation without mitral regurgitation in sheep. *Circulation* 1999; **100** (19 Suppl): II95–102.
 31. Boltwood CM, Tei C, Wong M, Shah PM. Quantitative echocardiography of the mitral complex in dilated cardiomyopathy: the mechanism of functional mitral regurgitation. *Circulation* 1983; **68**: 498–508.
 32. Levine RA, Handschumacher MD, Sanfilippo AJ, et al. Three-dimensional echocardiographic reconstruction of the mitral valve, with implications for the diagnosis of mitral valve prolapse. *Circulation* 1989; **80**: 589–98.
 33. Flachskampf FA, Chandra S, Gaddipati A, et al. Analysis of shape and motion of the mitral annulus in subjects with and without cardiomyopathy by echocardiographic 3-dimensional reconstruction. *J Am Soc Echocardiogr* 2000; **13**: 277–87.
 34. Gorman JH 3rd, Jackson BM, Enomoto Y, Gorman RC. The effect of regional ischemia on mitral valve annular saddle shape. *Ann Thorac Surg* 2004; **77**: 544–8.
 35. Matsui Y, Fukada Y, Suto Y, et al. Overlapping cardiac volume reduction operation. *J Thorac Cardiovasc Surg* 2002; **124**: 395–7.
 36. Matsui Y, Fukada Y, Naito Y, Sasaki S. Integrated overlapping ventriculoplasty combined with papillary muscle plication for severely dilated heart failure. *J Thorac Cardiovasc Surg* 2004; **127**: 1221–3.
 37. Suma H, Isomura T, Horii T, et al. Nontransplant cardiac surgery for end-stage cardiomyopathy. *J Thorac Cardiovasc Surg* 2000; **119**: 1233–44.
 38. Suma H, Isomura T, Horii T, Hisatomi K. Left ventriculoplasty for ischemic cardiomyopathy. *Eur J Cardiothorac Surg* 2001; **20**: 319–23.
 39. Ueno T, Sakata R, Iguro Y, Nagata T, Otsuji Y, Tei C. A new surgical approach to reduce tethering in ischemic mitral regurgitation by relocation of separate heads of posterior papillary muscle. *Ann Thorac Surg* 2005. (in press)
 40. Messas E, Guerrero JL, Handschumacher MD, et al. Chordal cutting: a new therapeutic approach for ischemic mitral regurgitation. *Circulation* 2001; **104**: 1958–63.
 41. Messas E, Pouzet B, Touchot B, et al. Efficacy of chordal cutting to relieve chronic persistent ischemic mitral regurgitation. *Circulation* 2003; **108** (Suppl 1): II111–5.
 42. Menicanti L, Di Donato M, Frigiola A, et al. Ischemic mitral regurgitation: intraventricular papillary muscle imbrication without mitral ring during left ventricular restoration. *J Thorac Cardiovasc Surg* 2002; **123**: 1041–50.
 43. Yamamoto H, Iguro Y, Sakata R, Arata K, Yotsumoto G. Effectively treating ischemic mitral regurgitation by chordal cutting in combination with ring annuloplasty and left ventricular reshaping approach. *J Thorac Cardiovasc Surg* 2005. (in press)
 44. Tibayan FA, Rodriguez F, Langer F, et al. Does septal-lateral annular cinching work for chronic ischemic mitral regurgitation? *J Thorac Cardiovasc Surg* 2004; **127**: 654–63.
 45. Hvass U, Tapia M, Baron F, Pouzet B, Shafy A. Papillary muscle sling: a new functional approach to mitral repair in patient with ischemic left ventricular dysfunction and functional mitral regurgitation. *Ann Thorac Surg* 2003; **75**: 809–11.